# GrowerFacts

Annuals Culture (revised 06/21/22)

# Dahlia Figaro<sup>™</sup> Series

(Dahlia x hybrida) Plug crop time: 4 to 5 weeks Transplant to finish: 5 to 6 weeks

• Premium pack item is ideal for borders and containers.

• Each variety contains several shades of that colour.

# **General Information**

Exposure	Bloom Season	Height	Spread	Spacing
Partial Sun	Spring, Late Spring, Summer, Late Summer, Autumn	14-18 in. (36-46 cm)	10-12 in. (25-30 cm)	10-14 in. (25-36 cm)

#### Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
RAW	288	1	4-5	3-7	5.5-5.8 pH 0.5-0.75 mmhos/cm	Yes

# Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 3	Level 3	Level 2-3	Level 2-3
Temperature	68-73°F (20-23°C)	68-73°F (20-23°C)	65-70°F (18-21°C)	65-70°F (18-21°C)
Light		1,500-3,000 f.c. (16,100-32,300 Lux)	1,500-3,000 f.c. (16,100-32,300 Lux)	2,500-4,000 f.c. (26,900-43,100 Lux)
Fertiliser		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)

#### **Propagation Key Tips**

If a germ chamber is used, move trays to the greenhouse at first sign of germination.

## **Growing on to Finish**

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 52-60°F (11-16°C)	5.8-6.2 pH	225 to 300 ppm N	Obligate Long Day
(night) 52-60°F (11-16°C)	0.75-1.0 mmhos/cm	(1.5 to 2.0 EC)	

#### **Daylength Notation**

Ideal daylength for Figaro Dahlia is in between 12 and 14 hours. Short days will cause tuber formation. Too long days (16 hours) will keep the plants more vegetative and delay flower initiation.

## **Crop Scheduling**

Container Size	Plugs/Pot	Crop Time	Season	PGR
306 Pack/1801	1 (ppp)	5-6 (weeks)	Spring	-
4"/4.5"/Quart/10 cm	1 (ppp)	5-6 (weeks)	Spring	-

#### **Chemical Sensitivity**

A-Rest, Bonzi, B-Nine/Alar and Cycocel can be used for plant control.

#### **Finishing Key Tips**

Very responsive to B-Nine/Alar. Also responsive to day/night temperature differential (DIF), and plants are shorter with a negative DIF.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

# **Variety Pictures**









ellow Shades

Orange Shades

Violet Shades

PanAmerican Seed.

PanAmerican Seed Co. 622 Town Road,West Chicago, Illinois, USA, 60185 +1 800-231-7065 PanAmSeed.com

<sup>™</sup> denotes a trademark of and <sup>®</sup> denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries. ©2024 Ball Horticultural Company